Asthma (Adults)

Asthma (Adults)

- •Most recent data (2014): 8% or about 122,600 King County adults age 18+.
- •Trend over time: 2000-14: flat
- •Description: Ever been told by a doctor, nurse or other health professional you have asthma and still have asthma
- •Healthy People 2020 Goal: None listed for this indicator.

Asthma (adults), King County, 2010-2014 average

	Percent	Lower Cl	Upper CI
King County (adults age 18+)	9	8	9
AGE			
18-24	7	6	10
25-44	9	8	10
45-64	9	8	10
65+	9	8	10
GENDER		·	
Vale	7	6	8
- emale	11	10	12
RACE/ETHNICITY			
AIAN	16 §	9 §	28 §
Asian	4	3	5
Black	10	7	14
lispanic	7	5	10
Multiple	11	8	16
NHPI	6 §	2 §	19 §
White	10	9	11
HOUSEHOLD INCOME			
\$15,000	15	12	19
\$15,000 to \$24,999	11	9	14
\$25,000 to \$34,999	11	8	14
\$35,000 to \$49,999	9	7	11
\$50,000 to \$74,999	7	6	8
\$75,000+	7	7	8
REGION		·	
East	8	7	9
North	8	6	10
Seattle	9	7	10
South	10	9	11
STATE			
Vashington State	10	9	10

Comparisons:

Higher than King County rate (yellow)
Lower than King County rate (green)

Notes:

Source: Behavioral Risk Factor Surveillance System.

Prepared by Public Health - Seattle & King County, APDE, 07/2016.


CI is 95% Confidence Interval.

 $\$ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

 $^{^{\}star}$ Too few cases to protect confidentiality and/or report reliable rates.

Asthma (adults) King County, 2010-2014 average



Source: Behavioral Risk Factor Surveillance System.
Prepared by Public Health - Seattle & King County, APDE, 07/2016.

|-----| Confidence interval shows range that includes true value 95% of the time.

* Too few cases to protect confidentiality and/or report reliable rates

* Too few cases to protect confidentiality and/or report reliable rates. § Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

Asthma (adults) by health reporting areas, King County, 2010-2014 average

	Percent	Lower CI	Upper CI
King County (adults age 18+)	9	8	9
Auburn	14	9	20
Aubum-North	11	7	17
Auburn-South	18	10	30
Bear Creek/Carnation/Duvall	8	5	11
Bellevue	7	5	9
Bellevue-Central	7 §	4 §	13 §
Bellewe-NE	7	4	12
Bellevue-South	7 §	4 §	13 §
Bellevue-West	5 §	2§	10 §
Black Diamond/Enumclaw/SE County	12	8	19
Bothell/Woodinville	10	5	16
Burien	12	8	18
Covington/Maple Valley	9	6	13
Des Moines/Normandy Park	8	5	14
East Federal Way	5 §	3 §	10 §
Fairwood	12 §	6 §	22 §
Federal Way	12 %	8	17
Fed Way-Central/Military Rd	12	9	20
	9 §		
Fed Way-Dash Point/Woodmont	<u> </u>	5 \$	17 §
Issaquah	12	7	20
Kenmore/LFP	6	4	10
Kent	8	6	11
Kent-East	11	6	18
Kent-SE	8	6	13
Kent-West	6 §	3 §	12 §
Kirkland	10	7	15
Kirkland	11	7	19
Kirkland North	9	6	14
Mercer Isle/Pt Cities	7	5	12
Newcastle/Four Creeks	8	5	13
North Highline	9 §	4 \$	17 §
Redmond	6	4	10
Renton	10	7	14
Renton-East	9 §	5 §	17 §
Renton-North	10	6	18
Renton-South	10	6	16
Sammamish	9	5	13
SeaTac/Tukwila	7	4	11
Seattle	9	7	10
Ballard	10	6	16
Beacon/Gtown/S.Park	12 §	6 §	21 §
Capitol Hill/E.lake	9	6	13
Central Seattle	11 §	6 §	19 [§]
Delridge	12	7	19
Downtown	8	5	13
Fremont/Greenlake	4	2	7
NE Seattle	6	4	10
North Seattle	9	6	13
NW Seattle	10	6	16
QA/Magnolia	8 §	4 §	15 [§]
SE Seattle	8	5	14
West Seattle	9	6	12
Shoreline	8	5	11
Snoqualmie/North Bend/Skykomish	9	5	14
Vashon Island	3 §	1 §	7 §

Comparisons:

Higher than King County rate (yellow)

Lower than King County rate (green)

Notes

Source: Behavioral Risk Factor Surveillance System.

Prepared by Public Health - Seattle & King County, APDE, 07/2016.


CI is 95% Confidence Interval.

§ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

 $^{^{\}star}$ Too few cases to protect confidentiality and/or report reliable rates.


Asthma (adults) by health reporting areas King County, 2010-2014 average



Note: HRA labels on the map match the chart on the right, listed in alphabetical order Source: Behavioral Risk Factor Surveillance System.

Prepared by Public Health - Seattle & King County, APDE, 07/2016.

Asthma (adults) King County, 2000-2014, three-year rolling averages



Trends over time: King County (2000-14: flat); East KC (2000-14: flat); North KC (2000-14: flat); Seattle (2000-14: flat); South KC (2000-14: flat); Washington State (2000-14: flat) Source: Behavioral Risk Factor Surveillance System.

Prepared by Public Health - Seattle & King County, APDE, 07/2016.

Produced by the

<u>Assessment, Policy Development & Evaluation Unit</u>

<u>Public Health - Seattle & King County</u>

07/16

